No.



8400063

THE DAIKHED STAYLES OF ANTERIOA

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Nickerson American Plant Breeders, Inc.

Wilherens. There has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OF ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF EIGHTEEN THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FILE AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCUED OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, UPORTING IT, OR EXPORTING IT, OR SUBJECT OF THE PLANT VARIETY PROTECTION ACT.

INITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS OFF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS THE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

WHEAT

'Success'

In Testimony Whereof, I have hereunto set my hand and caused the seal of the Plant Variety Erotection Office to be affixed at the City of Washington

this 21th day of September in the year of our Lord one thousand nine hundred and eighty-five.

Secretary of Agriculture

Lenxeth WEva. Commissioner Plant Variety Protection Office

Stant Variety Grotection Office Agricultural Marketing Service

AND XX AND XX I

	UNITED STATES DEPARTMEI	CETING SERVICE			FORM APPROVE, OMB NO. 46-F 2323
	APPLICATION FOR PLANT VARIE INSTRUCTIONS: See Reverse.				nt variety protection ನಾಜ್ಯ mpleted application ತೆಂಗಣ
	1a. TEMPORARY DESIGNATION OF	16. VARIETY NAM	E		AL USE ONLY
	HS79-400	Success		PV NUMBER 840	00063
v v	2. KIND NAME	3. GENUS AND SPE	CIES NAME	FILING DATE 3-8-84	TIME XXX
	Hard Red Spring Wheat	Triticum ae:	<u>stivum</u>	FEE RECEIVED	DATE
	Gramineae	5. DATE OF DETER 1)Spring o 2)Spring o	of '79	\$,800 \$	3-8-84
	6. NAME OF APPLICANT(S)	ļ	t and No. or R.F.D. No.,	City, State, and ZIP	8. TELEPHONE AREA
, es x x	NICKEESON) North American Plant Breeders	Code) 520	l Johnson Drive sion, KS 6620	, P.O. Box 2955	913-384-4940 KS 303-532-3721 CO
<i>نظا</i> هد م	9. IF THE NAMED APPLICANT IS NOT A PE ORGANIZATION: (Corporation, partnersh		10. IF INCORPORATE DATE OF INCORE	ED, GIVE STATE AND PORATION	11. DATE OF INCOR- PORATION
	Corporation		Delaware, Janua	•	1-19-83
	12. NAME AND MAILING ADDRESS OF APPI ALL PAPERS: G. E. Dixon R.E.M P.O. Box 2955 Mission, KS 66	LICANT REPRESENTA HEINER ROS 6201	ATIVE(S), IF ANY, TO S ERY F. BRITSE. He- P.O. Box Berthoud	iner or C. Brun 30	ATION AND RECERVE
2.2	13. CHECK BOX BELOW FOR EACH ATTACH		JC. 0044;	, 00 00010	
	X 13A. Exhibit A, Origin and Bree	ding History of the	Variety (See Section 5	2 of the Plant Variety	Protection Act.)
	X 13B. Exhibit B, Novelty Statem	ent.			
	X 13C. Exhibit C, Objective Descr	iption of the Variety	(Request form from	Plant Variety Protecti	on Office.)
	13D. Exhibit D, Additional Desc X 13E. Exhibit E. Qual	•	•	-	
:	14a. DOES THE APPLICANT(S) SPECIFY THAT SEED? (See Section 83(a). (If "Yes," answe			HETY NAME ONLY AS NO	A CLASS OF CERTIFIE
	14b. DOES THE APPLICANT(S) SPECIFY THAT LIMITED AS TO NUMBER OF GENERATI		14c. IF "YES," TO 14I TION BEYOND B		ATIONS OF PRODUC-
٠	X YES NO		X FOUNDATION		X CERTIFIED
	15a. DID THE APPLICANT(S) FILE FOR PROTI name of countries and dates.)	ECTION OF THIS VAR	RIETY IN OTHER COUP	NTRIES? TYES	X NO (If "Yes, " give
	15b. HAVE RIGHTS BEEN GRANTED THIS VA and dates.)	ARIETY IN OTHER CO	UNTRIES? YES	NO (If "Yes,"	give name of countries
	•				
•		·			
٠.	16. DOES THE APPLICANT(S) AGREE TO THE	E PUBLICATION OF H	IS/HER (THEIR) NAME	E(S) AND ADDRESS IN	THE OFFICIAL
•	17. The applicant(s) declare(s) that a viable replenished upon request in accordance	e sample of basic seed with such regulatio	d of this variety will b ns as may be applicabl	e furnished with the a	pplication and will be
	The undersigned applicant(s) is (are) the variety is distinct, uniform, and stable at 42 of the Plant Variety Act.	e owner(s) of this se	xually reproduced no	vel plant variety, and	
	Applicant(s) is (are) informed that false	e representation here	in can jeopardize prot	ection and result in p	enalties.
_	17 February 1984		_ Rober	tE Heiner	
	20 Color 1604			SIGNATURE OF APPLI	CANT)
-	(DATE)	No.	WI WA	SIGNATURE OF APPLI	CANT)
	FORM GR-470 (1-78)		1 -		

Exhibit A

Origin and Breeding History of Success

Pedigree: Era/3/Polk/Tobari//Fletcher

History:

The F2 population was obtained from the University of Minnesota in 1977. Success originated from a single F3 head selection in 1978. An F4 bulk of this selection was first entered into yield trials in 1979 under the experimental number HS79-400. This line has been in wide scale testing in the Red River Valley of the upper Midwest during the 1980 through 1983 seasons. Additionally, it was tested throughout the spring wheat region in the Uniform Regional Nursery during the 1982 season.

Two hundred ninety-six heads were pulled from the F5 generation of the original bulk in the 1980 season. These were grown in head-rows for initial purification at Berthoud, Colorado in 1981. Of these, 16 rows (5%) were discarded due to taller height and two rows (.7%) were discarded due to earlier heading. In the 1982 season, the remaining bulked head-rows were grown to produce the breeder seed lot.

Success is uniform and pure. Less than .5% of the plants have been rogued from the breeder seed field in 1982. Ninety-five percent of these rogued plants were three to five centimeters taller than Success. Less than .05% of these taller plants may be expected in subsequent generations.

Exhibit B

Novelty Statement

Success is most similar to the hard red spring wheat Kitt. However, it can be distinguished by the following morphological charateristics:

- Success and Kitt differ significantly in acuminate beak length; Success has a shorter beak length (see statistical data Exhibit E., page 2).
- Success is a green plant color at anthesis. Kitt is a blue-green plant color at anthesis.

Anova Table for Beak Length of Varieties Success and Kitt

Source	<u>df</u>	<u>ss</u>	<u>ms</u>
Total	49	290.55	
VAR	1	222.18	222.18
Error	48	68.37	1.42

F Test =
$$154.47**$$
 LSD (.05) = .67

<u>Variety</u>	Mean
Kitt	7.54 mm
Success	3.32 mm

The probability that the means for beak length are significantly different at the 5% and 1% alpha level.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK, MEAT, GRAIN & SEED DIVISION BELTSVILLE, MARYLAND 20705

EXHIBIT C (Wheat)

OBJECTIVE DESCRIPTION OF VARIETY

	TICUM SPP.)
NAME OF APPLICANTISI NICKERSON	FOR OFFICIAL USE ONLY
North American Plant Breeders Inc.	
ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code) 5201 Johnson Drive, P.O. Box 2955	VARIETY NAME OR TEMPORARY
Mission, KS 66201	DESIGNATION
111331011; 13 00201	Success
Place the appropriate number that describes the varietal character Place a zero in first box (c.s. 0 8 9 or 0 9) when number	
I. KIND:	
	= POLISH 6 = POULARD 7 = CLUB
2. TYPE:	1 = SOFT 3 = OTHER (Specify)
1 1 = SPRING 2 = WINTER 3 = OTHER (Specify)	2 2 = HARD
2 1 = WHITE 2 = RED 3 = OTHER (Specity)	
3. SEASON - NUMBER OF DAYS FROM TO:	
0 8 9 FIRST FLOWERING Planting	0 9 5 LAST FLOWERING
4. MATURITY (50% Flowering):	
NO. OF DAYS EARLIER THAN	1 = ARTHUR 2 = SCOUT 3 = CHRIS
0 1 NO. OF DAYS LATER THAN	4 = LEMHI 5 = NUGAINES 6 = LEEDS 7=Kitt
5. PLANT HEIGHT (From soil level to top of head):	
0 8 8 cm. HIGH	
0 5 cm. TALLER THAN	. 7
	1 = ARTHUR 2 = SCOUT 3 = CHRIS
CM. SHORTER THAN	4 = LEMHI 5 = NUGAINES 6 = LEEDS 7=Kitt
6. PLANT COLOR AT BOOTING (See reverse):	7. ANTHER COLOR:
2 1 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN	1 1 = YELLOW 2 = PURPLE
8. STEM:	
Anthocyanin: 1 = ABSENT 2 = PRESENT	Waxy bloom: 1 = ABSENT 2 = PRESENT
2 Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT	1 Internodes: 1 = HOLLOW 2 = SOLID
0 4 NO. OF NODES (Originating from node above ground)	1 8 CM. INTERNODE LENGTH BETWEEN FLAG LEAF AND LEAF BELOW
9. AURICLES:	
Anthocyanin: 1 = ABSENT 2 = PRESENT	2 Hairiness: 1 = ABSENT 2 = PRESENT
10. LEAF:	•
Flag leaf at 1 = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify):	Flag leaf: 1 = NOT TWISTED 2 = TWISTED
2 Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT	2 Waxy bloom of flag leaf sheath: 1 = ABSENT 2 = PRESENT
1 5 MM, LEAF WIDTH (First leaf below flag loaf)	2 7 CM. LEAF LENGTH (First leaf below flag leaf):

كالمناة والمناف المتاب والمساوية والمساوية والمساوية والمناوة والمناف والمناف المراوة والمراوة والمراوية والمراوية والمراوية		THE RESIDENCE OF THE PROPERTY		
Success			8400063	
FORM GR-470-6 (REVERSE	E)	·		<u>. </u>
11. HEAD:				
3 Density: 1 = LAX	^{2 = DENSE} 3 = middense average 48.0		ING 2 = STRAP 3 = CLAVATE (Specify)	
4 Awnedness: 1 = Aw		3 = AWNLETED 4 = AWNE	D	
Color at maturity:	= WHITE 2 = YELLOW 3 = PINK 4	= RED		
5	= BROWN 6 = BLACK 7 = OTHE	R (Specify):		
9.0 CM. LENGTH		1 1 мм. wютн		
12. GLUMES AT MATURI	TY:			
2 Length: 1 = SHORT 3 = LONG((CA. 7 mm.) 2 = MEDIUM (CA. 8 mm.) CA. 9 mm.) average 8.1 mm		N(CA. 3 mm.) 2 = MEDIUM (CA. 3.5 mm A. 4 mm.) average 3.5 mm	.)
· · · · · · · · · · · · · · · · · · ·				
• • • • • • • • • • • • • • • • • • •	ING 2 = OBLIQUE 3 = ROUNDED RE 5 = ELEVATED 6 = APICULATE	3 Beak: 1 = OBTUSE	2 = ACUTE 3 = ACUMINATE average 3.	4_ma
13. COLEOPTILE COLOR	•	14. SEEDLING ANTHOCY		
1 1 = WHITE 2 = RI	ED 3 = PURPLE	1 I = ABSENT 2	= PRESENT	
5. JUVENILE PLANT GR	OWTH HABIT:			
2 1 = PROSTRATE	2 = SEMI-ERECT 3 = EREC	:T		
I. SEED:			-	
3 Shape: I = OVATE	2 = OVAL 3 = ELLIPTICAL	Cheek: 1 = ROUND	ED 2 = ANGULAR	
2 Brush: 1 = SHORT			DLLARED 2 = COLLARED	
Phenol reaction (See instructions):	1 = IVORY • 2 = FAWN 3 = LT. BROWN 4 = BROWN 5 = BLACK 4=91% BY	n rown 5=9% Black		
3 Color: 1 = WHITE	2 = AMBER 3 = RED 4 = PURPLE	5 = OTHER (Specify)		
6.7 MM. LENGTH	3 7 MM. WIDTH	3 6 GM. PER 1000	SEEDS	
7. SEED CREASE:	- , Well-Man-	,		_
Width:] = 60% OR L	ESS OF KERNEL 'WINOKA'	Depth: 1 = 20% Of	R LESS OF KERNEL 'SCOUT'	
U 2 = 80% OR LE	ESS OF KERNEL 'CHRIS'	2 = 35% OF	LESS OF KERNEL 'CHRIS'	
	AS WIDE AS KERNEL "LEMHI"	3 = 50 % OF	LESS OF KERNEL 'LEMHI'	·· -
B. DISEASE: (0 = Not Test	red, 1 = Susceptible, 2 = Resistant) 3 =	moderately suscep	tible 4 = moderately	resis tar
2 (Races)	4 (Races) field races	0 STRIPE RUST	0 LOOSE SMUT	
O POWÖERY MILDEW	O BUNT	O OTHER (Specify)		
9. INSECT: (0 = Not Teste	od, 1 = Susceptible, 2 = Resistant) 3	= moderately susce	ptible 4 = moderately	_ resista
0 SAWFLY	O APHID (Bydv.)	O GREEN BUG	. O CEREAL LEAF BEETLE	
OTHER (Specify)	HESSIAN FLY	0 GP 0 A	0 B 0 C	
	RACES:	0 0 0 5	0 F 0 G	
INDICATE WINCO WASH	TV NOST OLOSEL V DESENDA DO TIVO	UDWITTED		<u>.</u>
CHARACTER	NAME OF VARIETY	UBMITTED: CHARACTER	NAME OF VARIETY	-
Plant tillering	Kitt	Seed size	Kitt	-
l enf size	Kitt	Seed shape	Kitt	-

INSTRUCTIONS

GENERAL: The following publications may be used as a reference aid for the standardization of terms and procedures for completing this form:

(a) L.W. Briggle and L. P. Reitz, 1963, Classification of Triticum Species and Wheat Varieties Grown in the United States, Technical Bulletin 1278, United States Department of Agriculture.

Coleoptile elongation

Seedling pigmentation

(b) W.E. Walls, 1965, A Standardized Phenol Method for Testing Wheat Seeds for Varietal Purity, contribution No. 28 to the handbook of seed testing prepared by the Association of Official Seed Analysts. (See attachment.)

Kitt

Kitt

Kitt

Kitt

Leaf color

Leaf carriage

Exhibit D

Additional Description of Success


NICKERSON

Success is a hard red spring wheat developed by North American Plant Breeders Inc. It was tested as the experimental number HS79-400.

Success is a tall semidwarf variety with medium-strong straw strength and late maturity. Its breadmaking quality is satisfactory. Its protein and bake absorption levels are slightly higher than the variety, Era.

Juvenile plant growth habit is semi-erect. Plant color at boot is green with a recurved, twisted flag leaf. Head shape is tapering to strap, middense, awned and head color is white at maturity. Glumes are of medium length and width with oblique shoulders and acumminate beaks. Seed shape is elliptical to ovate with rounded cheeks. Seed crease width is narrow and depth is shallow.

Success is adapted to the spring wheat region of North and South Dakota, and Minnesota.



Breeders
Plant
<u>=</u>
2
ican.
Amer
North

HARD RED SPRING WHEAT QUALITY

PAGE 1

				WHEA	WHEAT FLOUR	-	OUP ITY			_	BAKING	BAKING GUALITY	7					
YEAR	SAMPLE	PLE LOC	TEST OC WT.	PROT	FLR	FLR PROT	FLR ASH	MIX	ABS	MIX	DOUGH LOAF CHAR VOL	1	GRUMB.		덩	MILL	BAKE	TOTAL.
			16/Bu	14%mb	×	14%mb	14%mb	æ	×	eta	œ	S.	æ	æ	æ			
64	HS79-400		HU 57.6		72.7	11.4	0.493	٨	62.0	A. W.		1000+	~	9	6	23.0	83-B	158-C
70	HS79-400		å,		75.6		0.420	8 0	66.0	4.0	6 0	945	9	~	ത	81-B	86-B	167-B
် မ	HS79-400		4 10		63.1	٦.	0.377	4	58.0	3.0	9	1000	භ	~	6	G-09	73-0	135-D
1 8	HS79-4(CR 36.4	E	6 6 6 7	11.7	0.511	9	63.0	က : က :	6 0	973	ĸ.	N	c 0	67-D	86-B	133-0
T 8	HS79-400				0.83	···f	0.472	Φ)	65.0	e e	ø	066	æ	m	თ	84-B	92-A	176-B
-	HS79-400		.∀ 58.5		72.8	10.7	265.0	Ç	64.0	(T)	σ	040	ď		α	0.00 U-0.00	86.	1 8 4
े (\ व	HS79-400	2	0.00	14.3	9	13.0	0.495	œ	63.0	i i i	n or	1000+	σ	- σ	o o r	0-64	0-00	169-1
/ (A)	HS79-4		V 62.5		72.2	12.3	0.457	o co	63.0	, c	, et	1000) c) (٠ ٥	0 0	T 100	101
8	HS79-400				62.0	13.0	0.487	n	63.0	4	, O	830	σ	6	, Q1	0-89	83.4	123-0
2	HS79-400				75.4	12.3	0.481	9	63.0	2.8	8	840	69	8	o.	81-B	78-C	159-0
			•		•													
83	H579-4(73.2		0.427	9	64.0	3.0	80	940	Φ	100	ζn	81-B	86-B	167-B
83	HS79-400		HU 53.2	16.2	68.3	14.9	0.521	^	65.0	ອີ	80	1000+	69	0	6	74-C	93-4	167-B
	AVE	AVERAGE	58.2	14.0	70.5	12.5	0.478	Ģ	63.4	3.6	50)	963	€)	80	o	79-C	87-B	166-B
			•															
R	ERA	= 1	57.2		73.9		0.426	1 0 (63.0	4	N	920	9	9	On (84-B	82-B	166-B
£ 3	ERA	J 3		12.6	74.6 6.6.0	11.0	0.400	ໝ ∢	20.0	0 4	9 N	980	~ ¤	φ r	ന മ	7-6-K	7 d 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	132-0
3 =		. (٠.		70 A	σ	0 526	r	3	9 0		980	1	- 1	, a	- C9	0 E	144-5
	ERA	J	E 36.8	12.6	66.7	o.	0.563) 4 0	66.0	9) co	1000+	. eo	. 40	: o	78 7-87	93-P	171-B
							į	1	i (1							(1
18	ERA		0.09 7.09 7.09	12.6	, c , c	20.5	104.0	ភ ម	0.19	0 e	ם ת	2 G	pα	\ a	20 0	744	7 - SE	1.601
2 6	1001	ى ر	61.7) a		0.452	o co	93.0	4) O	200	۸ () ec	, 0) (1)	87-B	166-B
3 2	ERA) I			20.7		0.423	. 4	60.09	0	. co	1000+	. EQ	. \$1	, o	74-C	83-B	157-C
63	ERA	J	CX 60.8		73.1		0.447	မ	62.0	4.3	8	006	60	Ø	σ	79-C	84-B	163-B
												·						
83	ERA	U			72.2		0.486	m	62.0	4.0	80	1000+	8	ው	60	80-B	83-B	
83	ERA	.	HU 54.6	13.3	9.79	ᅻ.	205.0	ĸ	. 64.0	4.0	σ,	1000+	on.	Q	ው	74-C	92-A	166-B
	AVE	AVERAGE	59.1	13.7	21.0	12,1	0.468	ဖ	61.7	4. U	c	\$66	8	€	φ.	79-C	86-B	165-B
						1												
	ا المارية (TING 1 CONDING	1 PENT	\$ 1500°		10000V	U	ا ا	T-OFFECTTONABLE	L G	i L	אומריקם	TORE					
RERATINGS:		9-10-EXCELLENT	LENT	8=600D		7 MACCEPTABLE	19. E	3-6-008	5-6-0UESTIONABLE		1-4=0	1-4=UNACCEPTABLE	TABLE					-